

1. Helmholtz decomposition

11. October 2022

Exercise 1.

Consider following velocity vectors:

$$\mathbf{u}_1 = (y, 0, 0), \mathbf{u}_2 = (-y, x, 0), \mathbf{u}_3 = \left(-\frac{y}{x^2 + y^2}, \frac{x}{x^2 + y^2}, 0 \right)$$

Plot the velocity fields and compute their curl. In which sense does it hold that in the rotating flow, the velocity curl is non-zero?

Solution:

The fields are depicted in Fig. 1. Velocity \mathbf{u}_1 (locally) represents for example the situation of two parallel planar surfaces moving against each other in their planes.

Curl computations:


$$\nabla \times \mathbf{u}_1 = (0, 0, -1)$$

$$\nabla \times \mathbf{u}_2 = (0, 0, 2)$$

$$\nabla \times \mathbf{u}_3 = \left(0, 0, \frac{x^2 + y^2 - 2x^2}{(x^2 + y^2)^2} + \frac{x^2 + y^2 - 2y^2}{(x^2 + y^2)^2} \right) = (0, 0, 0)$$

Only the field \mathbf{u}_3 has therefore zero curl. On the other hand, vector fields \mathbf{u}_1 a \mathbf{u}_2 are vortical.

Continuum mechanics describes behaviour in individual points. The vector field \mathbf{u}_1 is vortical, because it causes "vortical" motion for example around the point $(0, 0, 0)$, similarly as the vector field \mathbf{u}_2 . On



Obrázek 1: Plots for Exercise 1.

the other hand, vector field \mathbf{u}_3 is irrotational, because the particles just "circle" around the point $(0, 0, 0)$.

Exercise 2.

Consider vector fields from exercise 1.

a) Are these fields divergent? b) If it holds that the curl of the given vector field is identically zero in \mathbb{R}^3 , can we deduce something about its divergence? (Think about the Helmholtz decomposition.)

Solution:

a) All the fields from exercise 1 have zero divergence.

b) Yes, in principle. From the Helmholtz decomposition, we know that

$$\mathbf{u} = \nabla\phi + \nabla \times \mathbf{A},$$

where

$$\phi(\mathbf{r}) = -\frac{1}{4\pi} \int_{\mathbb{R}^3} \frac{\nabla \cdot \mathbf{u}(\mathbf{r}')}{|\mathbf{r}' - \mathbf{r}|} dv', \tag{1}$$

$$\mathbf{A}(\mathbf{r}) = \frac{1}{4\pi} \int_{\mathbb{R}^3} \frac{\nabla \times \mathbf{u}(\mathbf{r}')}{|\mathbf{r}' - \mathbf{r}|} dv'. \tag{2}$$

This implies that if $\nabla \times \mathbf{u} \equiv 0$, then $\mathbf{A} \equiv \mathbf{0}$, and therefore also $\nabla \times \mathbf{A} \equiv \mathbf{0}$. If we consider nontrivial vector field \mathbf{u} , it must hold $\nabla\phi \neq \mathbf{0}$. However, according to Eq. (1), this cannot happen if $\nabla \cdot \mathbf{u} \equiv 0$. Therefore, if it holds for a vector field that both its divergence and curl are zero, the field must be, as a result of the Helmholtz decomposition, identically equal zero.

Why this doesn't hold for a function \mathbf{u}_3 from the first exercise: This field is not continuous enough - it is not defined in the origin, Helmholtz decomposition therefore cannot be applied. Helmholtz decomposition in \mathbb{R}^3 assumes that the field is at least C^2 (and decays to infinity at least as $1/r$).

Exercise 3.

For the velocity field $\mathbf{u} = (-y, x, 0)$, find an arbitrary scalar and vector potential ϕ and \mathbf{A} , so that

$$\mathbf{u} = \nabla\phi + \nabla \times \mathbf{A} \tag{3}$$

inside a bounded domain in \mathbb{R}^3 .

Solution:

Vector field \mathbf{u} has zero divergence. It holds

$$0 = \nabla \cdot \mathbf{u} = \nabla \cdot \nabla\phi + \nabla \cdot \nabla \times \mathbf{A} = \Delta\phi + 0.$$

Scalar potential ϕ must therefore fulfil the condition $\Delta\phi = 0$. There is a lot of such functions (we are working inside a domain of unspecified shape, so we aren't bothering with any boundary conditions). We can choose for example $\phi = x$. Then $\nabla\phi = (1, 0, 0)$ and from equation (3) follows

$$\nabla \times \mathbf{A} = \mathbf{u} - \nabla\phi = (-1 - y, x, 0).$$

We are therefore looking for a function $\mathbf{A} = (A^x, A^y, A^z)$ meeting the conditions

$$\begin{pmatrix} \partial_y A^z - \partial_z A^y \\ \partial_z A^x - \partial_x A^z \\ \partial_x A^y - \partial_y A^x \end{pmatrix} = \begin{pmatrix} -1 - y \\ x \\ 0 \end{pmatrix}.$$

Again, there is a lot of such vectors. We can choose for example $A^z = 0$. For the remaining components, we then get $A^x = xz$ and $A^y = (y + 1)z$.

Altogether, we've found a scalar potential $\phi = x$ and a vector potential $\mathbf{A} = (xz, (y + 1)z, 0)$ that can be used for the decomposition of the vector field \mathbf{u} .