

Knot-theoretic ternary groups

Maciej Niebrzydowski¹, Agata Pilitowska²
Anna Zamojska-Dzienio²

¹Institute of Mathematics, Faculty of Mathematics, Physics and Informatics,
University of Gdańsk

²Faculty of Mathematics and Information Science, Warsaw University of
Technology

SSAOS 2018

Ternary quasigroups

Ternary quasigroups

A **ternary groupoid** is a non-empty set A with a ternary operation $[\]: A^3 \rightarrow A$.

Ternary quasigroups

A **ternary groupoid** is a non-empty set A with a ternary operation $[\]: A^3 \rightarrow A$.

A ternary groupoid $(A, [\])$ is a **ternary quasigroup** if for every $a, b, c \in A$ each of the following equations is uniquely solvable for $z \in A$:

$$[zab] = c;$$

$$[azb] = c;$$

$$[abz] = c.$$

Ternary quasigroups

A **ternary groupoid** is a non-empty set A with a ternary operation $[\]: A^3 \rightarrow A$.

A ternary groupoid $(A, [\])$ is a **ternary quasigroup** if for every $a, b, c \in A$ each of the following equations is uniquely solvable for $z \in A$:

$$[zab] = c;$$

$$[azb] = c;$$

$$[abz] = c.$$

Each ternary quasigroup $(A, [\])$ has left, middle and right **cancellation property**, e.g. for all $x, y, a, b \in A$

$$[xab] = [yab] \Rightarrow x = y \text{ (left-cancellativity).}$$

Ternary groups

An operation $[\]: A^3 \rightarrow A$ is **associative** if for all $a, b, c, d, e \in A$

$$[[abc]de] = [a[bcd]e] = [ab[cde]].$$

Ternary groups

An operation $[\]: A^3 \rightarrow A$ is **associative** if for all $a, b, c, d, e \in A$

$$[[abc]de] = [a[bcd]e] = [ab[cde]].$$

An associative ternary quasigroup is a **ternary group**.

Ternary groups

An operation $[\]: A^3 \rightarrow A$ is **associative** if for all $a, b, c, d, e \in A$

$$[[abc]de] = [a[bcd]e] = [ab[cde]].$$

An associative ternary quasigroup is a **ternary group**.

Generalizing binary groups: Kasner 1904, Dörnte 1929, Lehmer 1932, Post 1940 (n-groups, polyadic groups)

Knot-theoretic ternary groups

Knot-theoretic ternary groups

A **knot-theoretic ternary group** is a ternary group $(A, [\,])$ which satisfies

$$\begin{aligned} [[abc]cd] &= [[ab[bcd]][bcd]d], \\ [ab[bcd]] &= [a[abc][[abc]cd]], \end{aligned}$$

for all $a, b, c, d \in A$.

Knot-theoretic ternary groups


A **knot-theoretic ternary group** is a ternary group $(A, [])$ which satisfies

$$\begin{aligned} [[abc]cd] &= [[ab[bcd]][bcd]d], \\ [ab[bcd]] &= [a[abc][[abc]cd]], \end{aligned}$$

for all $a, b, c, d \in A$.

Virtual knot theory (Kauffman 1999)

Third Reidemeister move (flat version)



Niebrzydowski 2014

Semi-commutativity

Each knot-theoretic ternary group $(A, [])$ is **semi-commutative**:

$$[abc] = [cba]$$

for every $a, b, c \in A$.

Each knot-theoretic ternary group $(A, [])$ is **semi-commutative**:

$$[abc] = [cba]$$

for every $a, b, c \in A$.

Semi-commutativity = lack of orientation

Each knot-theoretic ternary group $(A, [])$ is **semi-commutative**:


$$[abc] = [cba]$$

for every $a, b, c \in A$.

Semi-commutativity = lack of orientation

Theorem (Gładzek-Gleichgewicht, 1982)

A ternary group is semi-commutative if and only if it is entropic.



$$[[a_1 a_2 a_3][b_1 b_2 b_3][c_1 c_2 c_3]] = [[a_1 b_1 c_1][a_2 b_2 c_2][a_3 b_3 c_3]].$$

Skew element

Skew element

In a ternary quasigroup $(A, [])$, for every element $a \in A$, the unique solution of the equation

$$[aza] = a$$

is the **skew element to** a (denoted by \bar{a}).

Skew element

In a ternary quasigroup $(A, [])$, for every element $a \in A$, the unique solution of the equation

$$[aza] = a$$

is the **skew element to** a (denoted by \bar{a}).

Theorem (Borowiec-Dudek-Duplij, 2006)

If $(A, [])$ is a ternary group, then

$$[\bar{a}aa] = [a\bar{a}a] = [aa\bar{a}] = a$$

$$[ba\bar{a}] = [b\bar{a}a] = [a\bar{a}b] = [\bar{a}ab] = b$$

$$\overline{[abc]} = [\bar{c}\bar{b}\bar{a}]$$

$$\bar{\bar{a}} = a$$

for every $a, b, c \in A$.

Knot-theoretic ternary groups

Theorem

Let $(A, [\])$ be a ternary group. $(A, [\])$ is a knot-theoretic ternary group if and only if $(A, [\])$ is semi-commutative and satisfies

$$[abb] = \bar{a}$$

for every $a, b \in A$.

Theorem

Let $(A, [\])$ be a ternary group. $(A, [\])$ is a knot-theoretic ternary group if and only if $(A, [\])$ is semi-commutative and satisfies

$$[abb] = \bar{a}$$

for every $a, b \in A$.

In particular, in each knot-theoretic group

$$[ccc] = \bar{c},$$

for every $c \in A$.

Example

Example

Let $k > 1$ be a natural number, $(\mathbb{Z}_k, +)$ be a cyclic group and a be a fixed element of \mathbb{Z}_k .

Define on the set \mathbb{Z}_k the ternary operation

$$[xyz] = x - y + z + a \pmod{k}.$$

Then $(\mathbb{Z}_k, [])$ is a knot-theoretic ternary group if and only if $2a = 0 \pmod{k}$ in \mathbb{Z}_k .

For each even k , there are exactly two knot-theoretic groups constructed in this way:

- 1 idempotent one for $a = 0$,
- 2 non-idempotent for a being the element of order 2 in \mathbb{Z}_k .

Let $(A, [\])$ be a ternary groupoid. A binary groupoid $(A, *)$, where $x * y = [xay]$ for some fixed $a \in A$, is a **retract** of $(A, [\])$ (denoted by $\text{ret}_a(A, [\])$).

Let $(A, [\])$ be a ternary groupoid. A binary groupoid $(A, *)$, where $x * y = [xay]$ for some fixed $a \in A$, is a **retract** of $(A, [\])$ (denoted by $\text{ret}_a(A, [\])$).

A retract $\text{ret}_a(A, [\])$ of a semi-commutative ternary group is an abelian group with the neutral element \bar{a} and the inverse of $x \in A$ given by $\overline{[axa]}$.

Knot-theoretic ternary group

Theorem

Each knot-theoretic ternary group $(A, [])$ is determined by an abelian group $(A, +)$ and an element $a \in A$ of order one or two in $(A, +)$. Then for every $x, y, z \in A$

$$[xyz] = x - y + z + a \quad \text{and} \quad \bar{x} = x + a.$$

Theorem

Each knot-theoretic ternary group $(A, [])$ is determined by an abelian group $(A, +)$ and an element $a \in A$ of order one or two in $(A, +)$. Then for every $x, y, z \in A$

$$[xyz] = x - y + z + a \quad \text{and} \quad \bar{x} = x + a.$$

We denote the knot-theoretic ternary group $(A, [])$ described above by $\mathcal{T}((A, +), a)$. The group $(A, +)$ is the **associated group** of $(A, []) = \mathcal{T}((A, +), a)$.

$$\text{ret}_a(\mathcal{T}((A, +), a)) = (A, +).$$

Isomorphisms of KTTG

Theorem

Let $(A_1, []_1) = \mathcal{T}((A_1, +_1), a)$ and $(A_2, []_2) = \mathcal{T}((A_2, +_2), b)$ be two knot-theoretic ternary groups. Then the following statements are equivalent:

- ❶ *$(A_1, []_1)$ and $(A_2, []_2)$ are isomorphic;*
- ❷ *there exists a group isomorphism $h: (A_1, +_1) \rightarrow (A_2, +_2)$ such that $b = h(a)$.*

Theorem

Let $(A_1, []_1) = \mathcal{T}((A_1, +_1), a)$ and $(A_2, []_2) = \mathcal{T}((A_2, +_2), b)$ be two knot-theoretic ternary groups. Then the following statements are equivalent:

- ❶ *$(A_1, []_1)$ and $(A_2, []_2)$ are isomorphic;*
- ❷ *there exists a group isomorphism $h: (A_1, +_1) \rightarrow (A_2, +_2)$ such that $b = h(a)$.*

Enumeration of isomorphism classes of knot-theoretic ternary groups (up to 64 elements)

$$\begin{array}{c}
 [cda] \\
 || \\
 b \\
 [bcd] = a \quad c = [dab] \\
 d \\
 || \\
 [abc]
 \end{array}$$

$$\begin{array}{c}
 \langle cda \rangle \\
 || \\
 b \\
 \langle bcd \rangle = a \quad c = \langle dab \rangle \\
 d \\
 || \\
 \langle abc \rangle
 \end{array}$$

Compatible ternary groups

Compatible ternary groups

Two ternary groups $(A, [])$ and $(A, < >)$ are **compatible** if the following condition is satisfied:

$$[ab < bcd >] = < a < abc > [< abc > cd] >,$$

for any $a, b, c, d \in A$.

Compatible ternary groups


Two ternary groups $(A, [])$ and $(A, < >)$ are **compatible** if the following condition is satisfied:

$$[ab < bcd >] = < a < abc > [< abc > cd] >,$$

for any $a, b, c, d \in A$.

- 1 Any knot-theoretic ternary group is compatible with itself.
- 2 For a finite abelian binary group (of an even rank) $(A, +)$ with a neutral element 0 and x an element of order 2: two (non-isomorphic) knot-theoretic ternary groups:
 $(A, []) = \mathcal{T}((A, +), 0)$ and $(A, < >) = \mathcal{T}((A, +), x)$ are compatible.

Mixed Reidemeister type three move



Theorem


For a pair of compatible knot-theoretic ternary groups $(A, [])$ and $(A, < >)$, and a diagram D of a flat virtual link, the number of knot-theoretic ternary group colorings of D is not changed by the Reidemeister moves used for flat virtual links.

Theorem

For a pair of compatible knot-theoretic ternary groups $(A, [])$ and $(A, < >)$, and a diagram D of a flat virtual link, the number of knot-theoretic ternary group colorings of D is not changed by the Reidemeister moves used for flat virtual links.

Example: The flat-virtual Hopf link is distinguished from the unlink (two disjoint unknotted loops) by two-element ternary groups.

Example



Thank you for your attention!