

On the lattice of complete join-endomorphisms of a complete lattice

by Sándor Radeleczki, Math. Institute, Univ. of Miskolc (joint research with Kalle Kaarli, Tartu Univ.)

56-th. SSAOS, Spindlerův Mlyn, Sept. 4, 2018.

1. Preliminaries

1. Preliminaries

Let L be a complete lattice. A map $f: L \rightarrow L$ is called a **complete join-endomorphism**, if it preserves arbitrary joins, i.e.

$$f\left(\bigvee_{i \in I} x_i\right) = \bigvee_{i \in I} f(x_i), \text{ for all } x_i \in L \text{ and any index set } I.$$

In particular, $f(0) = 0$. It was proved by G. Grätzer and E.T. Schmidt [GS] that the complete join-endomorphisms of L also form a complete lattice $\text{End}_\vee(L)$. Here $f \vee g$ for $f, g \in \text{End}_\vee(L)$ is defined as a pointwise join.

1. Preliminaries

Let L be a complete lattice. A map $f: L \rightarrow L$ is called a **complete join-endomorphism**, if it preserves arbitrary joins, i.e.

$$f\left(\bigvee_{i \in I} x_i\right) = \bigvee_{i \in I} f(x_i), \text{ for all } x_i \in L \text{ and any index set } I.$$

In particular, $f(0) = 0$. It was proved by G. Grätzer and E.T. Schmidt [GS] that the complete join-endomorphisms of L also form a complete lattice $\text{End}_\vee(L)$. Here $f \vee g$ for $f, g \in \text{End}_\vee(L)$ is defined as a pointwise join. They also proved that for a lattice L with finite bounded chains,

- (T1) $\text{End}_\vee(L)$ is distributive if and only if L is distributive.
- (T2) If L is not distributive, then $\text{End}_\vee(L)$ is not even semimodular.

1. Preliminaries

Let L be a complete lattice. A map $f: L \rightarrow L$ is called a **complete join-endomorphism**, if it preserves arbitrary joins, i.e.

$$f\left(\bigvee_{i \in I} x_i\right) = \bigvee_{i \in I} f(x_i), \text{ for all } x_i \in L \text{ and any index set } I.$$

In particular, $f(0) = 0$. It was proved by G. Grätzer and E.T. Schmidt [GS] that the complete join-endomorphisms of L also form a complete lattice $\text{End}_\vee(L)$. Here $f \vee g$ for $f, g \in \text{End}_\vee(L)$ is defined as a pointwise join. They also proved that for a lattice L with finite bounded chains,

- (T1) $\text{End}_\vee(L)$ is distributive if and only if L is distributive.
- (T2) If L is not distributive, then $\text{End}_\vee(L)$ is not even semimodular.

S. Valentini(1994) have shown that

- (Thm3) The complete join-endomorphisms of L form a **quantale** with respect to the composition and the pointwise join.

Our aim here is to determine the **completely join and meet-irreducible elements** in the lattice $\text{End}_\vee(L)$, because these can be viewed as some "building stones" of a complete lattice.

Definition 1.

Definition 1.

A **quantale** is an algebraic structure $\mathbb{Q} = (L, \vee, \odot)$, such that (L, \leq) is a complete lattice (induced by \vee) and (L, \odot) is a semigroup satisfying

$$a \odot \left(\bigvee_{i \in I} b_i \right) = \bigvee_{i \in I} (a \odot b_i) \text{ and } \left(\bigvee_{i \in I} b_i \right) \odot a = \bigvee_{i \in I} (b_i \odot a).$$

for all $a \in L$ and $b_i \in L, i \in I$.

Definition 1.

A **quantale** is an algebraic structure $\mathbb{Q} = (L, \vee, \odot)$, such that (L, \leq) is a complete lattice (induced by \vee) and (L, \odot) is a semigroup satisfying

$$a \odot \left(\bigvee_{i \in I} b_i \right) = \bigvee_{i \in I} (a \odot b_i) \text{ and } \left(\bigvee_{i \in I} b_i \right) \odot a = \bigvee_{i \in I} (b_i \odot a).$$

for all $a \in L$ and $b_i \in L$, $i \in I$. \mathbb{Q} is called **commutative**, if \odot is commutative, and \mathbb{Q} is **unital**, whenever (L, \odot) is a monoid. A unital quantale in which the neutral element of \odot coincides to the greatest element 1 of the lattice L is called **integral**.

Definition 1.

A **quantale** is an algebraic structure $\mathbb{Q} = (L, \vee, \odot)$, such that (L, \leq) is a complete lattice (induced by \vee) and (L, \odot) is a semigroup satisfying

$$a \odot \left(\bigvee_{i \in I} b_i \right) = \bigvee_{i \in I} (a \odot b_i) \text{ and } \left(\bigvee_{i \in I} b_i \right) \odot a = \bigvee_{i \in I} (b_i \odot a).$$

for all $a \in L$ and $b_i \in L$, $i \in I$. \mathbb{Q} is called **commutative**, if \odot is commutative, and \mathbb{Q} is **unital**, whenever (L, \odot) is a monoid. A unital quantale in which the neutral element of \odot coincides to the greatest element 1 of the lattice L is called **integral**.

Examples

(a) *Frames* are commutative quantales in which \odot and the meet operation \wedge coincide.

Definition 1.

A **quantale** is an algebraic structure $\mathbb{Q} = (L, \vee, \odot)$, such that (L, \leq) is a complete lattice (induced by \vee) and (L, \odot) is a semigroup satisfying

$$a \odot \left(\bigvee_{i \in I} b_i \right) = \bigvee_{i \in I} (a \odot b_i) \text{ and } \left(\bigvee_{i \in I} b_i \right) \odot a = \bigvee_{i \in I} (b_i \odot a).$$

for all $a \in L$ and $b_i \in L$, $i \in I$. \mathbb{Q} is called **commutative**, if \odot is commutative, and \mathbb{Q} is **unital**, whenever (L, \odot) is a monoid. A unital quantale in which the neutral element of \odot coincides to the greatest element 1 of the lattice L is called **integral**.

Examples

(a) *Frames* are commutative quantales in which \odot and the meet operation \wedge coincide.

(b) *The two-sided ideals* of a ring $(R, +, \cdot)$ with unit form an integral quantale $(\mathcal{I}(R), \vee, \bullet)$, where $(\mathcal{I}(R), \vee, \cap)$ is the complete lattice of the ideals of R and \bullet is their usual multiplication.

Definition 1.

A **quantale** is an algebraic structure $\mathbb{Q} = (L, \vee, \odot)$, such that (L, \leq) is a complete lattice (induced by \vee) and (L, \odot) is a semigroup satisfying

$$a \odot \left(\bigvee_{i \in I} b_i \right) = \bigvee_{i \in I} (a \odot b_i) \text{ and } \left(\bigvee_{i \in I} b_i \right) \odot a = \bigvee_{i \in I} (b_i \odot a).$$

for all $a \in L$ and $b_i \in L$, $i \in I$. \mathbb{Q} is called **commutative**, if \odot is commutative, and \mathbb{Q} is **unital**, whenever (L, \odot) is a monoid. A unital quantale in which the neutral element of \odot coincides to the greatest element 1 of the lattice L is called **integral**.

Examples

(a) *Frames* are commutative quantales in which \odot and the meet operation \wedge coincide.

(b) *The two-sided ideals* of a ring $(R, +, \cdot)$ with unit form an integral quantale $(\mathcal{I}(R), \vee, \bullet)$, where $(\mathcal{I}(R), \vee, \cap)$ is the complete lattice of the ideals of R and \bullet is their usual multiplication.

(c) *For any complete lattice* L , $(\text{End}_{\vee}(L), \vee, \cdot)$ is a unital quantale (its unit is the identity mapping on L .)

The complete join-endomorphisms of lattice L are strongly related with the so-called ordered relations of L .

2. Valentini's ordered relations

The complete join-endomorphisms of lattice L are strongly related with the so-called ordered relations of L .

2. Valentini's ordered relations

Definition 2.

A **\leq -ordered relation** is a binary relation $R \subseteq L \times L$ such that

- (1) For any $u, x, y, z \in L$, if $u \leq x$, $(x, y) \in R$ and $y \leq z$, then $(u, z) \in R$;
- (2) For any $t \in L$, $A \subseteq L$, if $(a, t) \in R$ for all $a \in A$, then $(\bigvee A, t) \in R$;
- (3) For any $t \in L$, $A \subseteq L$, if $(t, a) \in R$ for all $a \in A$, then $(t, \bigwedge A) \in R$.

The complete join-endomorphisms of lattice L are strongly related with the so-called ordered relations of L .

2. Valentini's ordered relations

Definition 2.

A **\leq -ordered relation** is a binary relation $R \subseteq L \times L$ such that

- (1) For any $u, x, y, z \in L$, if $u \leq x$, $(x, y) \in R$ and $y \leq z$, then $(u, z) \in R$;
- (2) For any $t \in L$, $A \subseteq L$, if $(a, t) \in R$ for all $a \in A$, then $(\bigvee A, t) \in R$;
- (3) For any $t \in L$, $A \subseteq L$, if $(t, a) \in R$ for all $a \in A$, then $(t, \bigwedge A) \in R$.

Let us denote by $OR_{\leq}(L)$ the ordered relations on L . Notice, that the smallest ordered relation with respect to \subseteq is the "corner" relation

$$\Gamma(L) = \{(x, y) \in L^2 \mid x = 0 \text{ or } y = 1\}.$$

The complete join-endomorphisms of lattice L are strongly related with the so-called ordered relations of L .

2. Valentini's ordered relations

Definition 2.

A **\leq -ordered relation** is a binary relation $R \subseteq L \times L$ such that

- (1) For any $u, x, y, z \in L$, if $u \leq x$, $(x, y) \in R$ and $y \leq z$, then $(u, z) \in R$;
- (2) For any $t \in L$, $A \subseteq L$, if $(a, t) \in R$ for all $a \in A$, then $(\bigvee A, t) \in R$;
- (3) For any $t \in L$, $A \subseteq L$, if $(t, a) \in R$ for all $a \in A$, then $(t, \bigwedge A) \in R$.

Let us denote by $OR_{\leq}(L)$ the ordered relations on L . Notice, that the smallest ordered relation with respect to \subseteq is the "corner" relation

$$\Gamma(L) = \{(x, y) \in L^2 \mid x = 0 \text{ or } y = 1\}.$$

The following characterization of ordered relations was given by K. Kaarli and V. Kuchmei:

Proposition 1.

Let R be a binary relation on a complete lattice L . Then the following are equivalent:

- (1) R is a \leq -ordered relation;*
- (2) R is the universe of a complete subdirect square of L containing the "corner" element $(0, 1)$;*
- (3) R is a complete compatible binary relation on L containing $\Gamma(L)$.*

Proposition 1.

Let R be a binary relation on a complete lattice L . Then the following are equivalent:

- (1) R is a \leq -ordered relation;*
- (2) R is the universe of a complete subdirect square of L containing the "corner" element $(0, 1)$;*
- (3) R is a complete compatible binary relation on L containing $\Gamma(L)$.*

Valentini proved that

Proposition 2.

For any complete lattice L , $(OR_{\leq}(L), \cap, \circ)$ is a unital quantale, with the multiplicative identity \leq .

(Here operation " \circ " denotes the usual relational product.)

He also proved (though, using a different terminology) that

Proposition 1.

Let R be a binary relation on a complete lattice L . Then the following are equivalent:

- (1) R is a \leq -ordered relation;*
- (2) R is the universe of a complete subdirect square of L containing the "corner" element $(0, 1)$;*
- (3) R is a complete compatible binary relation on L containing $\Gamma(L)$.*

Valentini proved that

Proposition 2.

For any complete lattice L , $(OR_{\leq}(L), \cap, \circ)$ is a unital quantale, with the multiplicative identity \leq .

(Here operation " \circ " denotes the usual relational product.)

He also proved (though, using a different terminology) that

Proposition 3.

For any complete lattice L , the quantales $(OR_{\leq}(L), \cap, \circ)$ and $(\text{End}_{\vee}(L), \vee, \cdot)$ are dually isomorphic.

This **dual isomorphism** (anti-isomorphism) is established by the following pair a mappings (see [KR]):

This **dual isomorphism** (anti-isomorphism) is established by the following pair a mappings (see [KR]):

The mapping $\psi: \text{OR}_{\leq}(L) \rightarrow \text{End}_{\vee}(L)$, $\psi(R) = f^R$, where for each $R \in \text{OR}_{\leq}(L)$ and $x \in L$,

$$f^R(x) := \bigwedge \{z \in L \mid (x, z) \in R\},$$

and its inverse mapping

$$\psi^{-1}(f) = R^f := \{(x, y) \in L^2 \mid f(x) \leq y\}.$$

This **dual isomorphism** (anti-isomorphism) is established by the following pair of mappings (see [KR]):

The mapping $\psi: \text{OR}_{\leq}(L) \rightarrow \text{End}_{\vee}(L)$, $\psi(R) = f^R$, where for each $R \in \text{OR}_{\leq}(L)$ and $x \in L$,

$$f^R(x) := \bigwedge \{z \in L \mid (x, z) \in R\},$$

and its inverse mapping

$$\psi^{-1}(f) = R^f := \{(x, y) \in L^2 \mid f(x) \leq y\}.$$

We used this isomorphism and the lattice $\text{OR}_{\leq}(L)$ to determine the completely join and meet-irreducible elements in the lattice $\text{End}_{\vee}(L)$ (i.e. in the corresponding quantale $(\text{End}_{\vee}(L), \vee, \cdot)$).

This **dual isomorphism** (anti-isomorphism) is established by the following pair of mappings (see [KR]):

The mapping $\psi: \text{OR}_{\leq}(L) \rightarrow \text{End}_{\vee}(L)$, $\psi(R) = f^R$, where for each $R \in \text{OR}_{\leq}(L)$ and $x \in L$,

$$f^R(x) := \bigwedge \{z \in L \mid (x, z) \in R\},$$

and its inverse mapping

$$\psi^{-1}(f) = R^f := \{(x, y) \in L^2 \mid f(x) \leq y\}.$$

We used this isomorphism and the lattice $\text{OR}_{\leq}(L)$ to determine the completely join and meet-irreducible elements in the lattice $\text{End}_{\vee}(L)$ (i.e. in the corresponding quantale $(\text{End}_{\vee}(L), \vee, \cdot)$).

An element $p \in L \setminus \{0\}$ of a complete lattice L is called **completely join-irreducible** if for any system $X \subseteq L$ the equality $p = \bigvee X$ implies $p \in X$. The set of completely join-irreducible elements of L is denoted by $J(L)$. The **completely meet-irreducible elements** of L are defined dually, consisting of a set $M(L)$.

In what follows, for any elements a, b of the lattice L we will use the notations $\downarrow a = \{x \in L \mid x \leq a\}$ and $\uparrow b = \{x \in L \mid x \geq b\}$.

In what follows, for any elements a, b of the lattice L we will use the notations $\downarrow a = \{x \in L \mid x \leq a\}$ and $\uparrow b = \{x \in L \mid x \geq b\}$.

Theorem 1.

(i) *The completely join-irreducible elements of $\text{OR}_{\leq}(L)$ are exactly the relations*

$$\Gamma(L) \cup (\downarrow j \times \uparrow m) \subseteq L^2$$

where $j \in J(L)$ and $m \in M(L)$.

(ii) *The completely meet-irreducible elements of $\text{End}_{\vee}(L)$ are exactly the maps of the form (with $j \in J(L)$, $m \in M(L)$):*

$$f(x) = \begin{cases} 0, & \text{if } x = 0 \\ m, & \text{if } x \leq j, x \neq 0 \\ 1, & \text{otherwise} \end{cases}$$

In what follows, for any elements a, b of the lattice L we will use the notations $\downarrow a = \{x \in L \mid x \leq a\}$ and $\uparrow b = \{x \in L \mid x \geq b\}$.

Theorem 1.

(i) *The completely join-irreducible elements of $\text{OR}_{\leq}(L)$ are exactly the relations*

$$\Gamma(L) \cup (\downarrow j \times \uparrow m) \subseteq L^2$$

where $j \in J(L)$ and $m \in M(L)$.

(ii) *The completely meet-irreducible elements of $\text{End}_{\vee}(L)$ are exactly the maps of the form (with $j \in J(L)$, $m \in M(L)$):*

$$f(x) = \begin{cases} 0, & \text{if } x = 0 \\ m, & \text{if } x \leq j, x \neq 0 \\ 1, & \text{otherwise} \end{cases}$$

Concerning the join-irreducible elements of $\text{End}_{\vee}(L)$ we have obtained the following results:

Theorem 2.

(i) Let $p \in J(L)$ and $m \in M(L)$. Then any mapping $f: L \rightarrow L$ of the form

$$f(x) = \begin{cases} 0, & \text{if } x \leq m \\ p, & \text{otherwise} \end{cases}$$

is a completely join-irreducible element of $\text{End}_V(L)$.

Theorem 2.

(i) Let $p \in J(L)$ and $m \in M(L)$. Then any mapping $f: L \rightarrow L$ of the form

$$f(x) = \begin{cases} 0, & \text{if } x \leq m \\ p, & \text{otherwise} \end{cases}$$

is a completely join-irreducible element of $\text{End}_\vee(L)$.

(ii) If L is a finite distributive lattice, then the (completely) join-irreducible elements of $\text{End}_\vee(L)$ are exactly the mappings given in the above (i).

Theorem 2.

(i) Let $p \in J(L)$ and $m \in M(L)$. Then any mapping $f: L \rightarrow L$ of the form

$$f(x) = \begin{cases} 0, & \text{if } x \leq m \\ p, & \text{otherwise} \end{cases}$$

is a completely join-irreducible element of $\text{End}_\vee(L)$.

(ii) If L is a finite distributive lattice, then the (completely) join-irreducible elements of $\text{End}_\vee(L)$ are exactly the mappings given in the above (i).

A **complete tolerance** of a complete lattice L is a reflexive, symmetric relation T on L compatible with arbitrary suprema and infima, i.e. for any system of pairs $(a_i, b_i) \in T$, $i \in I$ we have

$$\left(\bigwedge_{i \in I} a_i, \bigwedge_{i \in I} b_i \right) \in T \text{ and } \left(\bigvee_{i \in I} a_i, \bigvee_{i \in I} b_i \right) \in T.$$

Theorem 2.

(i) Let $p \in J(L)$ and $m \in M(L)$. Then any mapping $f: L \rightarrow L$ of the form

$$f(x) = \begin{cases} 0, & \text{if } x \leq m \\ p, & \text{otherwise} \end{cases}$$

is a completely join-irreducible element of $\text{End}_\vee(L)$.

(ii) If L is a finite distributive lattice, then the (completely) join-irreducible elements of $\text{End}_\vee(L)$ are exactly the mappings given in the above (i).

A **complete tolerance** of a complete lattice L is a reflexive, symmetric relation T on L compatible with arbitrary suprema and infima, i.e. for any system of pairs $(a_i, b_i) \in T$, $i \in I$ we have

$$\left(\bigwedge_{i \in I} a_i, \bigwedge_{i \in I} b_i \right) \in T \text{ and } \left(\bigvee_{i \in I} a_i, \bigvee_{i \in I} b_i \right) \in T.$$

If in addition the above T is transitive, then it is called a *complete congruence* of L .

If $T \subseteq L^2$ is a complete tolerance of L , then for any $x \in L$ we define:
 $x_T := \bigwedge \{z \in L \mid (x, z) \in T\}$ and $x^T := \bigvee \{z \in L \mid (x, z) \in T\}$.

If $T \subseteq L^2$ is a complete tolerance of L , then for any $x \in L$ we define:
 $x_T := \bigwedge \{z \in L \mid (x, z) \in T\}$ and $x^T := \bigvee \{z \in L \mid (x, z) \in T\}$.

It is known that for any complete tolerance T of L the map $\lambda_T: x \mapsto x_T$, $x \in L$ is a decreasing complete \vee -endomorphism, and $\mu_T: x \mapsto x^T$, $x \in L$ is an increasing complete \wedge -endomorphism of the lattice L .

If $T \subseteq L^2$ is a complete tolerance of L , then for any $x \in L$ we define:
 $x_T := \bigwedge \{z \in L \mid (x, z) \in T\}$ and $x^T := \bigvee \{z \in L \mid (x, z) \in T\}$.

It is known that for any complete tolerance T of L the map $\lambda_T: x \mapsto x_T$, $x \in L$ is a decreasing complete \vee -endomorphism, and $\mu_T: x \mapsto x^T$, $x \in L$ is an increasing complete \wedge -endomorphism of the lattice L .

Notice that T is a **congruence** if and only if λ_T and μ_T are idempotent mappings (i.e. $(\lambda_T)^2 = \lambda_T$.)

If $T \subseteq L^2$ is a complete tolerance of L , then for any $x \in L$ we define:
 $x_T := \bigwedge \{z \in L \mid (x, z) \in T\}$ and $x^T := \bigvee \{z \in L \mid (x, z) \in T\}$.

It is known that for any complete tolerance T of L the map $\lambda_T: x \mapsto x_T$, $x \in L$ is a decreasing complete \vee -endomorphism, and $\mu_T: x \mapsto x^T$, $x \in L$ is an increasing complete \wedge -endomorphism of the lattice L .

Notice that T is a **congruence** if and only if λ_T and μ_T are idempotent mappings (i.e. $(\lambda_T)^2 = \lambda_T$.)

Definition 3.

- (a) A lattice L with 0 is called **pseudocomplemented** if for each $x \in L$ there exists an $x^* \in L$ such that for any $y \in L$, $y \wedge x = 0 \Leftrightarrow y \leq x^*$.
- (b) A bounded lattice L is called **double-pseudocomplemented** if both L , both its dual L^d are pseudocomplemented.

If $T \subseteq L^2$ is a complete tolerance of L , then for any $x \in L$ we define:
 $x_T := \bigwedge \{z \in L \mid (x, z) \in T\}$ and $x^T := \bigvee \{z \in L \mid (x, z) \in T\}$.

It is known that for any complete tolerance T of L the map $\lambda_T: x \mapsto x_T$, $x \in L$ is a decreasing complete \vee -endomorphism, and $\mu_T: x \mapsto x^T$, $x \in L$ is an increasing complete \wedge -endomorphism of the lattice L .

Notice that T is a **congruence** if and only if λ_T and μ_T are idempotent mappings (i.e. $(\lambda_T)^2 = \lambda_T$.)

Definition 3.

- (a) A lattice L with 0 is called **pseudocomplemented** if for each $x \in L$ there exists an $x^* \in L$ such that for any $y \in L$, $y \wedge x = 0 \Leftrightarrow y \leq x^*$.
- (b) A bounded lattice L is called **double-pseudocomplemented** if both L , both its dual L^d are pseudocomplemented.

Proposition 4.

Let L be a finite lattice and $\theta \subseteq L^2$ a congruence of it which is maximal with respect to the property $\theta^* \neq \Delta$. Then the mapping $\lambda_\theta: L \rightarrow L$, $\lambda_\theta(x) = x_\theta$, is a decreasing join-irreducible element of $\text{End}_\vee(L)$.

As an immediate consequence of the first theorem (Thm1.) we obtain

As an immediate consequence of the first theorem (Thm1.) we obtain

Corollary 1.

If L is a double-pseudocomplemented complete lattice, then $\text{OR}_{\leq}(L)$ is pseudocomplemented and $\text{End}_{\vee}(L)$ is dually pseudocomplemented.

As an immediate consequence of the first theorem (Thm1.) we obtain

Corollary 1.

If L is a double-pseudocomplemented complete lattice, then $\text{OR}_{\leq}(L)$ is pseudocomplemented and $\text{End}_{\vee}(L)$ is dually pseudocomplemented.

Remark

The form of the atoms and the dual atoms (coatoms) of $\text{OR}_{\leq}(L)$ and $\text{End}_{\vee}(L)$ can be easily deduced from Thm1 and Thm2. For instance, we obtain:

As an immediate consequence of the first theorem (Thm1.) we obtain

Corollary 1.

If L is a double-pseudocomplemented complete lattice, then $\text{OR}_{\leq}(L)$ is pseudocomplemented and $\text{End}_{\vee}(L)$ is dually pseudocomplemented.

Remark

The form of the atoms and the dual atoms (coatoms) of $\text{OR}_{\leq}(L)$ and $\text{End}_{\vee}(L)$ can be easily deduced from Thm1 and Thm2. For instance, we obtain:





Corollary 2.

The dual atoms of $\text{End}_{\vee}(L)$ are exactly the maps of the form:

$$f(x) = \begin{cases} 0, & \text{if } x = 0, \\ d, & \text{if } x = a, \\ 1, & \text{otherwise} \end{cases}$$

where a is an atom, and d is a dual atom of L .

-  T. S Blyth, M. F. Janowitz: *Residuation Theory*, Pergamon (1972).
-  I. Chajda and S. Radeleczki, 0-conditions and tolerance schemes, *Acta Mathematica Univ. Comenianae*, Vol. LXXII, **2** (2003), 177-184.
-  G. Czédli, E.K. Horváth and S. Radeleczki, On tolerance lattices of algebras in congruence modular varieties, *Acta Math. Hungarica*, **100** (1-2) (2003), 9-17.
-  G. Grätzer and E.T. Schmidt, The lattice of all join-endomorphisms of a lattice, *Proc. of the AMS*, Vol. 9, No. 5 , **10**/1 (1958), 722-726.
-  Janowitz, M. F.: Tolerances and congruences on lattices, *Czechoslovak Math. J.* **36**, (1986), 108-115
-  K. Kaarli and A. Pixley, *Polynomial completeness in algebraic systems*, CRC Press, Boca Raton, 2000.

-  K. Kaarli, V. Kuchmei, S.E. Schmidt: Sublattices of the direct product, *Algebra Universalis* **58** (2008), 85-95.
-  K. Kaarli, and S. Radeleczki: Representation of integral quantales by tolerances, *Algebra Universalis* **79**, No5. (2018), <https://doi.org/10.1007/s00012-018-0484-1>
-  C.J. Mulvey, *Quantales*, in Hazewinkel, Michiel: *Encyclopedia of Mathematics*, Springer, 2001.
-  S. Valentini: Representation theorems for quantales, *Math. Logic Q* **40** (1994), 182-190.

Thank You for your kind attention!

Thank You for your kind attention!