

Representations of Boolean lattices by annihilators in associative rings

Małgorzata Jastrzębska

Siedlce University of Natural Sciences and Humanities

Summer School on General Algebra and Ordered Sets
September, 2018

All rings are associative with identity $1 \neq 0$, unless otherwise stated.

Let R be a ring and let $I \subseteq R$ be a subgroup of $(R, +)$.

I is a **left ideal** — $\forall_{r \in R} \forall_{i \in I} \quad ri \in I \quad (RI \subseteq I)$

I is a **right ideal** — $\forall_{r \in R} \forall_{i \in I} \quad ir \in I \quad (IR \subseteq I)$

I is a **two-sided ideal (an ideal)** — I is a left ideal that is also a right ideal.

The fact that I is a two-sided ideal of R will be denoted by $I \triangleleft R$.

$\mathcal{I}_l(R)$ – the set of all left ideals in R

$\mathcal{I}_r(R)$ – the set of all right ideals in R

$\mathcal{I}(R)$ – the set of two-sided ideals in R

$$\mathcal{I}(R) = \mathcal{I}_l(R) \cap \mathcal{I}_r(R)$$

If R is commutative ring then $\mathcal{I}_l(R) = \mathcal{I}_r(R) = \mathcal{I}(R)$.

For every ring R the sets $\mathcal{I}_l(R), \mathcal{I}_r(R), \mathcal{I}(R)$ ordered by inclusion are lattices with operations:

$$I \vee J = I + J \quad \text{and} \quad I \wedge J = I \cap J. \quad (1)$$

All these lattices have the smallest element 0 and the largest element R , and all $\mathcal{I}_l(R), \mathcal{I}_r(R), \mathcal{I}(R)$ are **complete and modular**.

The lattice $\mathcal{I}(R)$ is a sublattice of $\mathcal{I}_l(R)$ and also a sublattice of $\mathcal{I}_r(R)$.

Division ring — a ring R in which lattices $\mathcal{I}_l(R)$ and $\mathcal{I}_r(R)$ are a two-element chains $\{0, R\}$ (ring in which every nonzero element a has a multiplicative inverse, i.e., an element x with $ax = xa = 1$).

Examples: fields, ring of quaternions

Simple ring — a ring R in which the lattice $\mathcal{I}(R)$ is a two-element chain $\{0, R\}$.

Examples: division rings, $M_n(K)$ a full ring of n -by- n matrices over field K

R. L. Blair, Ideal lattices and the structure of rings, Trans. Amer. Math. Soc. 75 (1953), 136-153.

Theorem

A ring R has a right ideal lattice $\mathcal{I}_r(R)$ which is a Boolean lattice if and only if R is isomorphic with a finite direct sum of division rings.

Theorem

- 1) A ring R has a two-sided ideal lattice $\mathcal{I}(R)$ which is a Boolean lattice if and only if R is isomorphic with a finite direct sum of simple rings.*
- 2) The lattice $\mathcal{I}(R)$ is a Boolean lattice if and only if $\mathcal{I}(R)$ is complemented lattice.*

Agreements

If $\emptyset \neq X \subseteq R$ is a subset, then let:

$l(X)$ be the left annihilator of X in R

$$l(X) = \{r \in R : \forall_{x \in X} rx = 0\},$$

$r(X)$ be the right annihilator of X in R

$$r(X) = \{r \in R : \forall_{x \in X} xr = 0\},$$

$\mathcal{A}_l(R)$ the set of all left annihilators in R ,

$\mathcal{A}_r(R)$ the set of all right annihilators in R ,

$$\mathcal{A}_l(R) \subseteq \mathcal{I}_l(R) \text{ and } \mathcal{A}_r(R) \subseteq \mathcal{I}_r(R)$$

$\mathcal{A}_l(R)$ ordered by inclusion is a complete lattice with operations:

$$I \vee J = l(r(I) \cap r(J)) \quad \text{and} \quad I \wedge J = I \cap J \quad \text{for} \quad I, J \in \mathcal{A}_l(R)$$

$\mathcal{A}_r(R)$ ordered by inclusion is a complete lattice with operations:

$$I \vee J = r(l(I) \cap l(J)) \quad \text{and} \quad I \wedge J = I \cap J \quad \text{for} \quad I, J \in \mathcal{A}_r(R).$$

Properties:

- 1) If $I \triangleleft R$ then also $l(I) \triangleleft R$ and $r(I) \triangleleft R$.
- 2) For any $X \subseteq R$ the set $l(r(X))$ is the smallest left annihilator which contain X .
- 3) If $I \triangleleft R$ then also $l(r(I)) \triangleleft R$.
- 4) Both of lattices $\mathcal{A}_l(R)$ and $\mathcal{A}_r(R)$ have the smallest element 0 and the largest element R .

Theorem (J. Krempa, M.J.)



For every finite lattice L there exist noncommutative rings R_1, R_2 and commutative ring R_3 such that:

- 1) L can be represented as a sublattice of a lattice $\mathcal{A}_l(R_1)$.*
- 2) L can be represented as a sublattice of a lattice $\mathcal{A}_r(R_2)$.*
- 3) L can be represented as a sublattice of a lattice $\mathcal{A}_l(R_3) = \mathcal{A}_r(R_3)$.*

Galois correspondence

For any ring R the mapping $\mathcal{A}_l(R) \rightarrow \mathcal{A}_r(R)$ given by $I \rightarrow r(I)$ is an anti-isomorphism of complete lattices. The inverse function is given by $J \rightarrow l(J)$ for $J \in \mathcal{A}_r(R)$.

$$l(r(I + J)) = l(r(I) \cap r(J))$$



The set of two-sided ideals which are left annihilators, denoted by $\mathcal{B}_l(R)$, is a sublattice of $\mathcal{A}_l(R)$.

The set of two-sided ideals which are right annihilators, denoted by $\mathcal{B}_r(R)$, is a sublattice of $\mathcal{A}_r(R)$.

Lattices of annihilators

$$\mathcal{B}_l(R)$$

$$\mathcal{B}_r(R)$$

$$\mathcal{A}_l(R)$$

$$\mathcal{A}_r(R)$$

For any ring R if $\mathcal{A}_l(R) = \mathcal{A}_r(R)$ (as subsets of 2^R) then $\mathcal{A}_l(R) = \mathcal{A}_r(R) = \mathcal{B}_l(R) = \mathcal{B}_r(R)$.

If R is commutative ring then $\mathcal{A}_l(R) = \mathcal{A}_r(R)$.

Statement

If $\mathcal{I}_l(R)$ is a Boolean lattice then $\mathcal{A}_l(R) = \mathcal{A}_r(R)$ is a Boolean lattice equal to $\mathcal{I}_l(R)$.

If $\mathcal{I}(R)$ is a Boolean lattice then $\mathcal{B}_l(R) = \mathcal{B}_r(R)$ is a Boolean lattice equal to $\mathcal{I}(R)$.

Example of a ring in which $\mathcal{A}_l(R) = \mathcal{A}_r(R)$ is Boolean lattice and this lattice is not equal to $\mathcal{I}_l(R)$

Let K be a field and let $R = K[x, y]/(xy)$ be a homomorphic image of the polynomial ring in commuting variables x, y . This ring is commutative hence $\mathcal{A}_l(R) = \mathcal{A}_r(R) = \mathcal{B}_l(R) = \mathcal{B}_r(R)$ and the set of annihilators is $\{0, R, (x), (y)\}$. For example $(x)^2$ is an ideal but not an annihilator in R .

An **element** $x \in R$ is said to be **nilpotent** if there exists a natural number n such that $x^n = 0$.

The **left (right) ideal** I in R is said to be **nilpotent** if there exists a natural number n such that $I^n = 0$.

A ring R is called **reduced** if it has no nonzero nilpotent elements.

Let R be a ring with no nonzero nilpotent ideals. Then for each ideal I of R $l(I) = r(I)$. This implies that $\mathcal{B}_l(R) = \mathcal{B}_r(R)$.

Theorem (cf. S. Steinberg, *Lattice ordered Rings and Modules*)

Let R be a ring with no nonzero nilpotent ideals. Then $\mathcal{B}_l(R)$ is a Boolean lattice.

If R is a reduced ring then for every $x \in R$ we have $l(x) = r(x)$. Hence $\mathcal{A}_l(R) = \mathcal{A}_r(R)$ and R has no nonzero nilpotent ideals.

From this and the above theorem follows

Corollary

If R is a reduced ring then $\mathcal{A}_l(R) = \mathcal{A}_r(R)$ is a complete Boolean lattice.