

On minimality of generating sets of aggregation clones

Radomír Halaš, Zbyněk Kurač, Jozef Pócs



Palacký University
Olomouc

SSAOS 2018, Špindlerův Mlýn, Czech Republic
6th September 2018

1. Generating of an aggregation clone
2. Properties of functions χ and \oplus
3. The forms of a generating set

1. Generating of an aggregation clone

Aggregation function


Definition

Let (L, \leq) be a finite lattice and $k \in \mathbb{N}$. A mapping $A : L^k \rightarrow L$ is called a *k-ary aggregation function* on lattice L , if it is non-decreasing, i.e., for any $\mathbf{x}, \mathbf{y} \in L^n$:

$$\mathbf{x} \leq \mathbf{y} \Rightarrow A(\mathbf{x}) \leq A(\mathbf{y})$$

and it fulfills boundary conditions

$$A(0, \dots, 0) = 0 \quad \text{and} \quad A(1, \dots, 1) = 1.$$




Aggregation clone

Definition

Let \mathcal{O}_A be a set of all (finitary) operations defined on a set A . Then subset $\mathcal{C} \subseteq \mathcal{O}_A$ is called a *clone* on A if it contains all projection operations on A and is closed under the composition of functions.

- all aggregation functions form a clone (*aggregation clone* AGG_A)



The generating set of an aggregation clone using functions χ and \oplus

Definition

Let L be a finite lattice. Then functions $\chi_a : L \rightarrow L$ and $\oplus_a : L^2 \rightarrow L$ for all $a \in L$ are defined as follows

$$\chi_a(x) = \begin{cases} 1 & ; x \geq a, x \neq 0 \\ 0 & ; \text{otherwise,} \end{cases} \quad x \oplus_a y = \begin{cases} 1 & ; x = 1, y = 1 \\ 0 & ; x = 0, y = 0 \\ a & ; \text{otherwise.} \end{cases}$$

- the number of functions is $2n + 2$ for n -element lattice L

The generating set of an aggregation clone using functions χ and \oplus

Theorem

Let L be a finite lattice. Then the aggregation clone \mathcal{AGG}_L on L is generated by the lattice operations \vee, \wedge and by functions $\chi_a, \oplus_a, a \in L$.

- Let $A : L^k \rightarrow L$ be an aggregation function on lattice L . Then

$$A(\mathbf{x}) = \bigvee_{\mathbf{a} \in L_*^k} \left(\bigwedge_{i \in J_{\mathbf{a}}} \chi_{a_i}(x_i) \wedge \bigoplus_{i=1}^k \oplus_{A(\mathbf{a})} x_i \right)$$

where $\mathbf{x} = (x_1, \dots, x_k)$, $\mathbf{a} = (a_1, \dots, a_k)$,

$$L_*^k = L^k \setminus \{(0, \dots, 0), (1, \dots, 1)\}, J_{\mathbf{a}} = \{1 \leq i \leq k \mid a_i \neq 0\},$$

$$\bigoplus_{i=1}^k \oplus_{A(\mathbf{a})} x_i = x_1 \oplus_{A(\mathbf{a})} x_2 \oplus_{A(\mathbf{a})} \dots \oplus_{A(\mathbf{a})} x_k.$$

2. Properties of functions χ and \oplus

Properties of functions \oplus

Lemma

Let L be a finite lattice. Then the equalities are valid:

- i) $x \oplus_0 y = \chi_1(x) \wedge \chi_1(y),$
- ii) $x \oplus_1 y = \chi_0(x) \vee \chi_0(y).$

Moreover:

- i) $x \oplus_0 x = \chi_1(x),$
- ii) $x \oplus_1 x = \chi_0(x).$

Lemma

Let L be a finite lattice. Then the following equalities are valid for all $a, b \in L$:

- i) $(x \oplus_a y) \wedge (x \oplus_b y) = x \oplus_{a \wedge b} y,$
- ii) $(x \oplus_a y) \vee (x \oplus_b y) = x \oplus_{a \vee b} y.$



Properties of functions \oplus

Proposition

Let L be a finite lattice and let H_L be a minimal generating set of lattice L . Denote

$$G = \left\{ \chi_a \mid a \in L \right\} \cup \left\{ \oplus_b \mid b \in H_L \setminus \{0, 1\} \right\} \cup \left\{ \vee, \wedge \right\}.$$

Then the set $G_c = G \setminus \{\oplus_c\}$ does not generate the aggregation clone \mathcal{AGG}_L for any $c \in H_L \setminus \{0, 1\}$.



Properties of functions χ

Lemma

Let L be an n -element lattice, $n \geq 4$. Then for any $a, b, c \in L$ such that $a, b \neq 1$, $a \not\leq b$, $c \neq 0$ we have

- i) $\chi_0(x) = \chi_c(x \oplus_c x)$,
- ii) $\chi_1(x) = \chi_a(x \oplus_b x)$,
- iii) $\chi_0(x) = \chi_\alpha(x)$ if α is a unique atom of L .

Lemma

Let L be a finite lattice. Then for any $a_1, \dots, a_p \in L$ we have

- i) $\chi_{a_1}(x) \wedge \dots \wedge \chi_{a_p}(x) = \chi_{a_1 \vee \dots \vee a_p}(x)$,
- ii) $\chi_{a_1}(x) \vee \dots \vee \chi_{a_p}(x) = \chi_{a_1 \wedge \dots \wedge a_p}(x) = \chi_0(x)$ assuming a_1, \dots, a_p are just all the atoms of L .

Properties of atomary functions χ

- An atom $\alpha \Rightarrow \chi_\alpha \dots$ an *atomary function*

- $$\underbrace{\overbrace{\chi_{\alpha_1}(x)}^0 \vee \dots \vee \chi_{\alpha_i}(x) \vee \dots \vee \overbrace{\chi_{\alpha_q}(x)}^0}_{\Downarrow} = \chi_0(x), \quad x \mapsto x \wedge (x \oplus_{\alpha_i} x)$$

Lemma

Let L be a finite lattice with atoms $\alpha_1, \dots, \alpha_q$. Then all atomary functions χ_{α_i} can be generated according to the following formula:

$$\chi_{\alpha_i}(x) = \chi_0(x \wedge (x \oplus_{\alpha_i} x)).$$

3. The forms of a generating set

The cardinality of generating sets

- $\chi_{a_1}(x) \wedge \dots \wedge \chi_{a_p}(x) = \chi_{a_1 \vee \dots \vee a_p}(x) \Rightarrow J_L \dots$ join-irreducible elements of L
 $\chi_{\alpha_1}(x) \vee \dots \vee \chi_{\alpha_q}(x) = \chi_{\alpha_1 \wedge \dots \wedge \alpha_q}(x) \Rightarrow At_L \dots$ atoms of L
- Duality:
 $\mu_{a_1}(x) \vee \dots \vee \mu_{a_p}(x) = \mu_{a_1 \wedge \dots \wedge a_p}(x) \Rightarrow M_L \dots$ meet-irreducible elements of L
 $\mu_{\omega_1}(x) \wedge \dots \wedge \mu_{\omega_r}(x) = \mu_{\omega_1 \vee \dots \vee \omega_r}(x) \Rightarrow CoAt_L \dots$ coatoms of L
- $$\left. \begin{aligned} (x \oplus_a y) \wedge (x \oplus_b y) &= x \oplus_{a \wedge b} y \\ (x \oplus_a y) \vee (x \oplus_b y) &= x \oplus_{a \vee b} y \end{aligned} \right\} \Rightarrow H_L \dots \text{a generating set of lattice } L$$


Theorem

Let L be a finite lattice with q atoms and r coatoms. Let $H_L \subset L$ be a set of minimal cardinality generating the lattice L . Let $G \subset \mathcal{O}_A$ be a set of minimal cardinality generating the aggregation clone AGG_L . Then

$$|G| \leq \min(|J_L| - q, |M_L| - r) + |H_L| + 2.$$

The cardinality of (χ, \oplus) -generating sets


Example



- $G_1 = \{\chi_b, \chi_c, \chi_d\} \cup \{\oplus_a, \oplus_b, \oplus_c, \oplus_d\} \cup \{\vee, \wedge\} \dots$ greatest cardinality
- $G_2 = \{\chi_c, \chi_d\} \cup \{\oplus_a, \oplus_b, \oplus_c, \oplus_d\} \cup \{\vee, \wedge\}$
- $G_3 = \{\chi_c, \chi_d\} \cup \{\oplus_a, \oplus_b, \oplus_c, \oplus_d\} \cup \{\vee, \wedge\}$
- $G_4 = \{\chi_d\} \cup \{\oplus_a, \oplus_b, \oplus_c, \oplus_d\} \cup \{\vee, \wedge\}$

The cardinality of (χ, \oplus) -generating sets


Example



- $G_5 = \{\chi_b\} \cup \{\oplus_a, \oplus_c, \oplus_d\} \cup \{\vee, \wedge\}$
- $G_6 = \{\chi_d\} \cup \{\oplus_a, \oplus_c, \oplus_d\} \cup \{\vee, \wedge\}$
- $G_7 = \{\chi_c\} \cup \{\oplus_a, \oplus_b, \oplus_c\} \cup \{\vee, \wedge\}$
- $G_8 = \{\chi_d\} \cup \{\oplus_a, \oplus_b, \oplus_d\} \cup \{\vee, \wedge\}$
- $G_9 = \{\oplus_0, \oplus_a, \oplus_b, \oplus_c\} \cup \{\vee, \wedge\}$
- $G_{10} = \{\oplus_a, \oplus_b, \oplus_c, \oplus_1\} \cup \{\vee, \wedge\}$
- $G_{11} = \{\oplus_a, \oplus_b, \oplus_c, \oplus_d\} \cup \{\vee, \wedge\}$

The cardinality of (χ, \oplus) -generating sets

Example



- $G_{12} = \{\chi_d\} \cup \{\oplus_b, \oplus_c\} \cup \{\vee, \wedge\} \dots$ *smallest cardinality*
- $G_{13} = \{\oplus_b, \oplus_c, \oplus_d\} \cup \{\vee, \wedge\} \dots$ *smallest cardinality*
- $G_{14} = \{\oplus_a, \oplus_b, \oplus_c\} \cup \{\vee, \wedge\} \dots$ *smallest cardinality*
- $G_{15} = \{\oplus_a, \oplus_b, \oplus_c\} \cup \{\vee, \wedge\} \dots$ *smallest cardinality*

The greatest cardinality of a (χ, \oplus) -generating set

Proposition

Let $n \geq 4$ be a positive integer and L be n -element lattice. Let $\alpha \in L$ be an atom of lattice L . Consider the set

$$G = \left\{ \chi_b \mid b \in L \setminus \{0, \alpha, 1\} \right\} \cup \left\{ \oplus_c \mid c \in L \setminus \{0, 1\} \right\} \cup \left\{ \vee, \wedge \right\}.$$

Then the lattice L is a chain if and only if the set G is a minimal (χ, \oplus) -generating set of the aggregation clone \mathcal{AGG}_L .

- $|G| = 2n - 3$
- $n = 2 \quad \dots \quad G = \{\vee, \wedge\}$
- $n = 3 \quad \dots \quad G = \{\oplus_0, \oplus_a, \oplus_1\} \cup \{\vee, \wedge\}$

A generating set of an aggregation clone on an atomary lattice

Definition

Let L be a lattice. Let J_L be the set of join-irreducible elements of L and At_L be the set of atoms of L . Then L is called an *atomary lattice* if $J_L = At_L$.

Proposition

Let L be a finite atomary lattice. Let H_L be a minimal generating set of lattice L . Then the set $\{\oplus_b \mid b \in H_L\} \cup \{\vee, \wedge\}$ is a minimal (χ, \oplus) -generating set of the aggregation clone \mathcal{AGG}_L .

The smallest cardinality of a (χ, \oplus) -generating set

Hypothesis


Let L be an n -element atomary (resp. coatomary) lattice such that for another n -element atomary (resp. coatomary) lattice M we have

$$|H_L| \leq |H_M|,$$


where H_L and H_M are sets of minimal cardinality generating lattices L and M , respectively. Then the sets G_L and G_M of minimal cardinality generating aggregation clones \mathcal{AGG}_L and \mathcal{AGG}_M fulfill

$$|G_L| \leq |G_M|.$$

- $|G_L| = |H_L| + 2$






Example



- $G = \{\chi_n, \chi_k\} \cup \{\oplus_a, \oplus_c, \oplus_f, \oplus_h, \oplus_i, \oplus_m, \oplus_p\} \cup \{\vee, \wedge\}$

References

-  Botur M., Halaš R., Mesiar R., Pócs J. : *On generating of idempotent aggregations functions on bounded lattices*, Information Sciences 430-431 (2018) 39-45.
-  Halaš R., Pócs J. : *On the clone of aggregation functions on bounded lattices*, Information Sciences 329 (2016) 381–389.
-  Halaš R., Mesiar R., Pócs J. : *Generators of Aggregation Functions and Fuzzy Connectives*, IEEE Transactions on Fuzzy Systems 24(6) (2016) 1690-1694.

Thank you for your attention